

****TITLE****

*ASP Conference Series, Vol. **VOLUME**, **PUBLICATION YEAR***

****EDITORS****

Optical Spectroscopy of T dwarfs

Adam J. Burgasser

California Institute of Technology, MSC 103-33, Pasadena, CA 91125

J. Davy Kirkpatrick

Infrared Processing and Analysis Center, MSC 100-22, Pasadena, CA 91125

Abstract. Optical spectra are presented for a sample of T dwarfs, obtained using the Keck 10m Low Resolution Imaging Spectrograph. Dominant features are discussed, including possible temperature diagnostics.

We have used the Low Resolution Imaging Spectrograph (LRIS; Oke et al. 1995), mounted on the Keck 10m Telescope, to obtain red optical spectra of nine T dwarfs selected from Burgasser et al. (1999, 2000a, 2000c, 2001), Strauss et al. (1999), and Tsvetanov et al. (2000). Observations of 2MASS 1237+65, SDSS 1346-00, and SDSS 1624+00 are presented in Burgasser et al. (2000b), which discusses instrumental setup and data reduction; the remaining targets, discussed here, were observed on 5 March 2000 (UT).

Reduced spectra are shown in Figure 1a, normalized at 9200 Å and vertically offset by 0.5. The rapid rise in flux from 8000 to 10000 Å is likely due to the extremely pressure-broadened K I doublet at 7665 and 7699 Å, as noted in SDSS 1624+00 by Liebert et al. (2000). A slight rise in flux between 6500 and 7800 Å is seen in a few objects, most clearly in 2MASS 0559-14 and SDSS 1624+00 (Figure 1b), and is due to an opacity window between the K I doublet and the similarly broadened Na I D lines (5890 and 5896 Å; Reid et al. 2000). We note that the increase in flux on the red side of the K I doublet is greater in Gl 570D ($T_{eff} \sim 750$ K) than in 2MASS 0559-14 ($T_{eff} \sim 1200$ K). H₂O absorption at 9200 Å similarly increases from 2MASS 0559-14 to Gl 570D. The behavior of the Cs I lines (8521 and 8943 Å) is unclear due to poor signal-to-noise in these faint dwarfs, as well as the influence of CH₄ at 8900 Å; similarly, Rb I (7948 Å) is detected only in the high signal-to-noise spectrum of 2MASS 0559-14. These alkali features are seen to strengthen throughout the L dwarf sequence (Kirkpatrick et al. 1999). FeH (9896 Å), which weakens in mid- to late-L dwarfs, is seen in 2MASS 0559-14, SDSS 1624+00, and 2MASS 1225-27, but is not readily apparent in the other objects. Finally, the H α emission reported in 2MASS 1237+65 (Burgasser et al. 2000b) is not seen in any of the other T dwarfs, in agreement with activity relations observed in late-M and L dwarfs (Gizis et al. 2000), and further evidence that 2MASS 1237+65 may have an unusual activity mechanism. We propose that FeH, H₂O, and the red slope of the K I doublet provide potential temperature diagnostics in the optical regime.




Figure 1. Optical spectra of T dwarfs. (a) Full 6300–10100 Å spectra, with dominant features indicated. (b) 6300–8000 Å region, magnified 20× to show flux between Na I and K I doublets.

References

- Burgasser, A. J., et al. 1999, *ApJ*, 522, L65
 Burgasser, A. J., et al. 2000a, *ApJ*, 531, L57
 Burgasser, A. J., Kirkpatrick, J. D., Reid, I. N., Liebert, J., Gizis, J. E., & Brown, M. E. 2000b, *AJ*, 120, 473
 Burgasser, A. J., et al. 2000c, *AJ*, 120, 1100
 Burgasser, A. J., et al. 2001, *ApJ*, in prep.
 Gizis, J. E., Monet, D. G., Reid, I. N., Kirkpatrick, J. D., Liebert, J., & Williams, R. J. 2000, *AJ*, 120, 1085
 Kirkpatrick, J. D., et al. 1999, *ApJ*, 519, 802
 Liebert, J., Reid, I. N., Burrows, A., Burgasser, A. J., Kirkpatrick, J. D., & Gizis, J. E. 2000, *ApJ*, 533, L155
 Oke, J. B., et al. 1995, *PASP*, 107, 375
 Reid, I. N., Kirkpatrick, J. D., Gizis, J. E., Dahn, C. C., Monet, D. G., Williams, R. J., Liebert, J., & Burgasser, A. J. 2000, *AJ*, 119, 369
 Strauss, M. A., et al. 1999, *ApJ*, 522, L61
 Tsvetanov, Z. I., et al. 2000, *ApJ*, 531, L61